



NTNU

Norwegian University of Science and Technology

# Fenchel Duality Theory and a Primal-Dual Algorithm on Riemannian Manifolds

Ronny Bergmann

joint work with

R. Herzog, M. Silva Louzeiro, D. Tenbrinck, J. Vidal-Núñez.


IMADA Seminar, SDU, Odense,

June 2nd, 2022

# Manifold-valued Signal & Image Processing

Tasks in **image processing** are often phrased as an optimisation problem. **Here.** The pixel take values on a manifold

- ▶ phase-valued data ( $\mathbb{S}^1$ )
- ▶ wind-fields, GPS ( $\mathbb{S}^2$ )
- ▶ DT-MRI ( $\mathcal{P}(3)$ )
- ▶ EBSD, (grain) orientations ( $SO(n)$ )




Artificial noisy phase-valued data.

**Tasks.** Denoising, Inpainting, labeling (classification), deblurring,...

# Manifold-valued Signal & Image Processing

Tasks in **image processing** are often phrased as an optimisation problem.  
**Here.** The pixel take values on a manifold

- ▶ phase-valued data ( $\mathbb{S}^1$ )
- ▶ wind-fields, GPS ( $\mathbb{S}^2$ )
- ▶ DT-MRI ( $\mathcal{P}(3)$ )
- ▶ EBSD, (grain) orientations ( $SO(n)$ )



InSAR-Data of Mt. Vesuvius.

[Rocca, Prati, and Guarnieri 1997]

**Tasks.** Denoising, Inpainting, labeling (classification), deblurring,...

# Manifold-valued Signal & Image Processing

Tasks in **image processing** are often phrased as an optimisation problem. **Here.** The pixel take values on a manifold

- ▶ phase-valued data ( $\mathbb{S}^1$ )
- ▶ wind-fields, GPS ( $\mathbb{S}^2$ )
- ▶ DT-MRI ( $\mathcal{P}(3)$ )
- ▶ EBSD, (grain) orientations ( $SO(n)$ )




Artificial noisy data on the sphere  $\mathbb{S}^2$ .

**Tasks.** Denoising, Inpainting, labeling (classification), deblurring,...

# Manifold-valued Signal & Image Processing

Tasks in **image processing** are often phrased as an optimisation problem.  
**Here.** The pixel take values on a manifold

- ▶ phase-valued data ( $\mathbb{S}^1$ )
- ▶ wind-fields, GPS ( $\mathbb{S}^2$ )
- ▶ DT-MRI ( $\mathcal{P}(3)$ )
- ▶ EBSD, (grain) orientations ( $SO(n)$ )




Artificial diffusion data,  
 each pixel is a symmetric positive matrix.

**Tasks.** Denoising, Inpainting, labeling (classification), deblurring,...

# Manifold-valued Signal & Image Processing

Tasks in **image processing** are often phrased as an optimisation problem.  
**Here.** The pixel take values on a manifold

- ▶ phase-valued data ( $\mathbb{S}^1$ )
- ▶ wind-fields, GPS ( $\mathbb{S}^2$ )
- ▶ DT-MRI ( $\mathcal{P}(3)$ )
- ▶ EBSD, (grain) orientations ( $SO(n)$ )



DT-MRI of the human brain.

Camino Project: [cmic.cs.ucl.ac.uk/camino](http://cmic.cs.ucl.ac.uk/camino)

**Tasks.** Denoising, Inpainting, labeling (classification), deblurring,...

# Manifold-valued Signal & Image Processing

Tasks in **image processing** are often phrased as an optimisation problem.

**Here.** The pixel take values on a manifold

- ▶ phase-valued data ( $\mathbb{S}^1$ )
- ▶ wind-fields, GPS ( $\mathbb{S}^2$ )
- ▶ DT-MRI ( $\mathcal{P}(3)$ )
- ▶ EBSD, (grain) orientations ( $SO(n)$ )



Grain orientations in EBSD data.


MTEX toolbox: [mtex-toolbox.github.io](https://github.com/mTEX-toolbox)

**Tasks.** Denoising, Inpainting, labeling (classification), deblurring,...

# A $d$ -dimensional Riemannian manifold $\mathcal{M}$

## Notation.

- ▶ Geodesic  $\gamma(\cdot; p, q)$
- ▶ Tangent space  $\mathcal{T}_p\mathcal{M}$
- ▶ inner product  $(\cdot, \cdot)_p$
- ▶ Logarithmic map  $\log_p q = \dot{\gamma}(0; p, q)$
- ▶ Exponential map  $\exp_p X = \gamma_{p,X}(1)$   
where  $\gamma_{p,X}(0) = p$  and  $\dot{\gamma}_{p,X}(0) = X$
- ▶ Parallel transport  $P_{q \leftarrow p} Y$  "move"  
tangent vectors from  $\mathcal{T}_p\mathcal{M}$  to  $\mathcal{T}_q\mathcal{M}$



# The Model

We consider a minimization problem

$$\arg \min_{p \in \mathcal{C}} F(p) + G(\Lambda(p))$$

- ▶  $\mathcal{M}, \mathcal{N}$  are (high-dimensional) Riemannian Manifolds
- ▶  $F: \mathcal{M} \rightarrow \overline{\mathbb{R}}$  nonsmooth, (locally, geodesically) convex
- ▶  $G: \mathcal{N} \rightarrow \overline{\mathbb{R}}$  nonsmooth, (locally) convex
- ▶  $\Lambda: \mathcal{M} \rightarrow \mathcal{N}$  nonlinear
- ▶  $\mathcal{C} \subset \mathcal{M}$  strongly geodesically convex.

In image processing.

choose a model, such that finding a minimizer yields the reconstruction

# Splitting Methods & Algorithms

On a Riemannian manifold  $\mathcal{M}$  we have

- ▶ Cyclic Proximal Point Algorithm (CPPA) [Bačák 2014]
- ▶ (parallel) Douglas–Rachford Algorithm (PDRA) [RB, Persch, and Steidl 2016]

On  $\mathbb{R}^n$  PDRA is known to be equivalent to [Setzer 2011; O'Connor and Vandenberghe 2018]

- ▶ Primal-Dual Hybrid Gradient Algorithm (PDHGA) [Esser, Zhang, and Chan 2010]
- ▶ Chambolle-Pock Algorithm (CPA) [Chambolle and Pock 2011; Pock, Cremers, Bischof, and Chambolle 2009]

But on a Riemannian manifold  $\mathcal{M}$ :  no duality theory!

## Goals of this talk.

Formulate Duality on a Manifold

Derive a Riemannian Chambolle–Pock Algorithm (RCPA)

# Musical Isomorphisms

[Lee 2003]

The dual space  $\mathcal{T}_p^*\mathcal{M}$  of a tangent space  $\mathcal{T}_p\mathcal{M}$  is called **cotangent space**. We denote by  $\langle \cdot, \cdot \rangle$  the duality pairing.

We define the **musical isomorphisms**

- ▶  $b: \mathcal{T}_p\mathcal{M} \ni X \mapsto X^\flat \in \mathcal{T}_p^*\mathcal{M}$  via  $\langle X^\flat, Y \rangle = (X, Y)_p$  for all  $Y \in \mathcal{T}_p\mathcal{M}$
- ▶  $\sharp: \mathcal{T}_p^*\mathcal{M} \ni \xi \mapsto \xi^\sharp \in \mathcal{T}_p\mathcal{M}$  via  $(\xi^\sharp, Y)_p = \langle \xi, Y \rangle$  for all  $Y \in \mathcal{T}_p\mathcal{M}$ .

which introduces an inner product and parallel transport on/between  $\mathcal{T}_p^*\mathcal{M}$

## (Geodesic) Convexity

[Sakai 1996; Udriște 1994]

A set  $\mathcal{C} \subset \mathcal{M}$  is called (strongly geodesically) **convex** if for all  $p, q \in \mathcal{C}$  the geodesic  $\gamma(\cdot; p, q)$  is unique and lies in  $\mathcal{C}$ .

A function  $F: \mathcal{C} \rightarrow \overline{\mathbb{R}}$  is called (geodesically) **convex** if for all  $p, q \in \mathcal{C}$  the composition  $F(\gamma(t; p, q)), t \in [0, 1]$ , is convex.

# The Subdifferential

The **subdifferential** of  $F$  at  $p \in \mathcal{C}$  is given by

[Lee 2003; Udriște 1994]

$$\partial_{\mathcal{M}} F(p) := \{ \xi \in \mathcal{T}_p^* \mathcal{M} \mid F(q) \geq F(p) + \langle \xi, \log_p q \rangle \text{ for } q \in \mathcal{C} \},$$

where

- ▶  $\mathcal{T}_p^* \mathcal{M}$  is the dual space of  $\mathcal{T}_p \mathcal{M}$ ,
- ▶  $\langle \cdot, \cdot \rangle$  denotes the duality pairing on  $\mathcal{T}_p^* \mathcal{M} \times \mathcal{T}_p \mathcal{M}$

# The Euclidean Fenchel Conjugate

Let  $f: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  be proper and convex.

We define the **Fenchel conjugate**  $f^*: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  of  $f$  by



$$f^*(\xi) := \sup_{x \in \mathbb{R}^n} \langle \xi, x \rangle - f(x) = \sup_{x \in \mathbb{R}^n} \begin{pmatrix} \xi \\ -1 \end{pmatrix}^T \begin{pmatrix} x \\ f(x) \end{pmatrix}$$

- ▶ interpretation: maximize the distance of  $\xi^T x$  to  $f$
- ⇒ extremum seeking problem on the epigraph

The Fenchel **biconjugate** reads

$$f^{**}(x) = (f^*)^*(x) = \sup_{\xi \in \mathbb{R}^n} \langle \xi, x \rangle - f^*(\xi).$$

# Illustration of the Fenchel Conjugate



# Properties of the Euclidean Fenchel Conjugate

[Rockafellar 1970]

- ▶ The Fenchel conjugate  $f^*$  is **convex** (even if  $f$  is not)
- ▶  $f^{**}$  is the largest convex, lsc function with  $f^{**} \leq f$
- ▶ If  $f(x) \leq g(x)$  holds for all  $x \in \mathbb{R}^n$  then  $f^*(\xi) \geq g^*(\xi)$  holds for all  $\xi \in \mathbb{R}^n$
- ▶ **Fenchel–Moreau theorem**:  $f$  convex, proper, lsc  $\Rightarrow f^{**} = f$ .
- ▶ **Fenchel–Young inequality**:

$$f(x) + f^*(\xi) \geq \xi^T x \quad \text{for all } x, \xi \in \mathbb{R}^n$$

- ▶ For a proper, convex function  $f$

$$\xi \in \partial f(x) \Leftrightarrow f(x) + f^*(\xi) = \xi^T x$$

- ▶ For a proper, convex, lsc function  $f$ , then

$$\xi \in \partial f(x) \Leftrightarrow x \in \partial f^*(\xi)$$

# The Riemannian $m$ -Fenchel Conjugate

[RB, Herzog, Silva Louzeiro, Tenbrinck, and Vidal-Núñez 2021]

alternative approaches: [Ahmadi Kakavandi and Amini 2010; Silva Louzeiro, RB, and Herzog 2022]

**Idea:** Introduce a point on  $\mathcal{M}$  to “act as” 0.

Let  $m \in \mathcal{C} \subset \mathcal{M}$  be given and  $F: \mathcal{C} \rightarrow \overline{\mathbb{R}}$ .

The  $m$ -Fenchel conjugate  $F_m^*: \mathcal{T}_m^* \mathcal{M} \rightarrow \overline{\mathbb{R}}$  is defined by

$$F_m^*(\xi_m) := \sup_{X \in \mathcal{L}_{\mathcal{C}, m}} \{ \langle \xi_m, X \rangle - F(\exp_m X) \},$$

where  $\mathcal{L}_{\mathcal{C}, m} := \{X \in \mathcal{T}_m \mathcal{M} \mid q = \exp_m X \in \mathcal{C} \text{ and } \|X\|_p = d(q, p)\}$ .

Let  $m' \in \mathcal{C}$ . The  $mm'$ -Fenchel-biconjugate  $F_{mm'}^{**}: \mathcal{C} \rightarrow \overline{\mathbb{R}}$  is given by

$$F_{mm'}^{**}(p) = \sup_{\xi_{m'} \in \mathcal{T}_{m'}^* \mathcal{M}} \{ \langle \xi_{m'}, \log_{m'} p \rangle - F_m^*(P_{m \leftarrow m'} \xi_{m'}) \}.$$

usually we only use the case  $m = m'$ .

# Properties of the $m$ -Fenchel Conjugate

[RB, Herzog, Silva Louzeiro, Tenbrinck, and Vidal-Núñez 2021]

- ▶  $F_m^*$  is convex on  $\mathcal{T}_m^*\mathcal{M}$
- ▶  $F(p) \leq G(p)$  holds for all  $p \in \mathcal{C} \Rightarrow F_m^*(\xi_m) \geq G_m^*(\xi_m)$  holds for all  $\xi_m \in \mathcal{T}_m^*\mathcal{M}$
- ▶ Fenchel-Moreau theorem:  $F \circ \exp_m$  convex (on  $\mathcal{T}_m\mathcal{M}$ ), proper, lsc, then  $F_{mm}^{**} = F$  on  $\mathcal{C}$ .
- ▶ Fenchel-Young inequality: For a proper, convex function  $F \circ \exp_m$

$$\xi_p \in \partial_{\mathcal{M}} F(p) \Leftrightarrow F(p) + F_m^*(P_{m \leftarrow p} \xi_p) = \langle P_{m \leftarrow p} \xi_p, \log_m p \rangle.$$

- ▶ For a proper, convex, lsc function  $F \circ \exp_m$

$$\xi_p \in \partial_{\mathcal{M}} F(p) \Leftrightarrow \log_m p \in \partial F_m^*(P_{m \leftarrow p} \xi_p).$$

# Proximal Map

For  $F: \mathcal{M} \rightarrow \overline{\mathbb{R}}$  and  $\lambda > 0$  we define the **Proximal Map** as  
[Moreau 1965; Rockafellar 1970; Ferreira and Oliveira 2002]

$$\text{prox}_{\lambda F} p := \arg \min_{u \in \mathcal{M}} d(u, p)^2 + \lambda F(u).$$

- ! For a Minimizer  $u^*$  of  $F$  we have  $\text{prox}_{\lambda F} u^* = u^*$ .
- ▶ For  $F$  proper, convex, lsc:
  - ▶ the proximal map is unique.
  - ▶ **Proximal-Point-Algorithm:**  
 $x_k = \text{prox}_{\lambda F} x_{k-1}$  converges to  $\arg \min F$
- ▶  $q = \text{prox}_{\lambda F} p$  is equivalent to

$$\frac{1}{\lambda} (\log_q p)^b \in \partial_{\mathcal{M}} F(q)$$

# The Chambolle-Pock Algorithm

[Chambolle and Pock 2011]

From the pair of primal-dual problems

$$\begin{aligned} \min_{x \in \mathbb{R}^n} f(x) + g(Kx), \quad K \text{ linear,} \\ \max_{\xi \in \mathbb{R}^m} -f^*(-K^*\xi) - g^*(\xi) \end{aligned}$$

we obtain for  $f, g$  proper convex, lsc the optimality conditions (OC) for a solution  $(\hat{x}, \hat{\xi})$  as ,

**Chambolle–Pock Algorithm.** with  $\sigma > 0, \tau > 0, \theta \in \mathbb{R}$  reads

$$\begin{aligned} \partial f &\ni -K^*\hat{\xi} \\ \partial g^*(\hat{\xi}) &\ni K\hat{x} \\ \bar{\xi}^{(k+1)} &= \xi^{(k+1)} + \theta(\xi^{(k+1)} - \xi^{(k)}) \end{aligned}$$

## Saddle Point Formulation

Let  $F$  be geodesically convex,  $G \circ \exp_n$  be convex (on  $\mathcal{T}_n\mathcal{N}$ ).

From

$$\min_{p \in \mathcal{C}} F(p) + G(\Lambda(p))$$

we derive the saddle point formulation for the  $n$ -Fenchel conjugate of  $G$  as

$$\min_{p \in \mathcal{C}} \max_{\xi_n \in \mathcal{T}_n^*\mathcal{N}} \langle \xi_n, \log_n \Lambda(p) \rangle + F(p) - G_n^*(\xi_n).$$

But  $\Lambda: \mathcal{M} \rightarrow \mathcal{N}$  is a non-linear operator!

For Optimality Conditions and the Dual Problem: What's  $\Lambda^*$ ?

**Approach.** Linearization:

[Valkonen 2014]

$$\Lambda(p) \approx \exp_{\Lambda(m)} D\Lambda(m)[\log_m p]$$

# The Exact Riemannian Chambolle–Pock Algorithm (eRCPA)

[RB, Herzog, Silva Louzeiro, Tenbrinck, and Vidal-Núñez 2021]

**Input:**  $m, p^{(0)} \in \mathbb{R}^d$  ,  $n = \Lambda(m)$ ,  $\xi_n^{(0)} \in \mathbb{R}^d$  , and parameters  $\sigma, \tau, \theta > 0$

1:  $k \leftarrow 0$

2:  $\bar{p}^{(0)} \leftarrow p^{(0)}$

3: **while** not converged **do**

4:  $\xi_n^{(k+1)} \leftarrow \text{prox}_{\tau G_n^*}(\xi_n^{(k)} + \tau (\log_n \Lambda(\bar{p}^{(k)}))^b)$

5:  $p^{(k+1)} \leftarrow \text{prox}_{\sigma F} \left( p^{(k)} + P_{p^{(k)} \leftarrow m} \left( -\sigma D\Lambda(m)^* [\xi_n^{(k+1)}]^\# \right) \right)$

6:  $\bar{p}^{(k+1)} \leftarrow p^{(k+1)} + \theta(p^{(k+1)} - p^{(k)})$

7:  $k \leftarrow k + 1$

8: **end while**

**Output:**  $p^{(k)}$

# Generalizations & Variants of the RCPA

Classically

[Chambolle and Pock 2011]

- ▶ change  $\sigma = \sigma_k, \tau = \tau_k, \theta = \theta_k$  during the iterations
- ▶ introduce an acceleration  $\gamma$
- ▶ relax dual  $\bar{\xi}$  instead of primal  $\bar{p}$  (switches lines 4 and 5)

Furthermore we

[RB, Herzog, Silva Louzeiro, Tenbrinck, and Vidal-Núñez 2021]

- ▶ introduce the **IRCPA**: linearize  $\Lambda$ , i. e., adopt the Euclidean case from

[Valkonen 2014]

$$\log_n \Lambda(\bar{p}^{(k)}) \rightarrow P_{n \leftarrow \Lambda(m)} D\Lambda(m) [\log_m \bar{p}^{(k)}]$$

- ▶ choose  $n \neq \Lambda(m)$  introduces a parallel transport

$$D\Lambda(m)^* [\xi_n^{(k+1)}] \rightarrow D\Lambda(m)^* [P_{\Lambda(m) \leftarrow n} \xi_n^{(k+1)}]$$

- ▶ change  $m = m^{(k)}, n = n^{(k)}$  during the iterations

## A Constant and a Conjecture

We define

$$C(k) := \frac{1}{\sigma} d^2(p^{(k)}, \tilde{p}^{(k)}) + \langle \bar{\xi}_n^{(k)}, D\Lambda(m)[\zeta_k] \rangle,$$

where

$$\zeta_k = P_{m \leftarrow p^{(k)}}(\log_{p^{(k)}} p^{(k+1)}) - P_{p^{(k)} \leftarrow \tilde{p}^{(k)}}(\log_{\tilde{p}^{(k)}} \hat{p}) - \log_m p^{(k+1)} + \log_m \hat{p},$$

and  $\hat{p}$  is a minimizer of the primal problem.

### Remark.

For  $\mathcal{M} = \mathbb{R}^d$ :  $\zeta_k = \tilde{p}^{(k)} - p^{(k)} = -\sigma(D\Lambda(m))^*[\bar{\xi}_n^{(k)}] \Rightarrow C(k) = 0$ .

### Conjecture.

Assume  $\sigma\tau < \|D\Lambda(m)\|^2$ . Then  $C(k) \geq 0$  for all  $k > K$ ,  $K \in \mathbb{N}$ .

# Convergence of the IRCPA

## Theorem.

[RB, Herzog, Silva Louzeiro, Tenbrinck, and Vidal-Núñez 2021]

Let  $\mathcal{M}$ ,  $\mathcal{N}$  be Hadamard. Assume that the linearized problem

$$\min_{p \in \mathcal{M}} \max_{\xi_n \in \mathcal{T}_n^* \mathcal{N}} \langle (D\Lambda(m))^*[\xi_n], \log_m p \rangle + F(p) - G_n^*(\xi_n).$$

has a saddle point  $(\hat{p}, \hat{\xi}_n)$ .

Choose  $\sigma, \tau$  such that

$$\sigma\tau < \|D\Lambda(m)\|^2$$

and assume that  $C(k) \geq 0$  for all  $k > K$ . Then it holds

1. the sequence  $(p^{(k)}, \xi_n^{(k)})$  remains bounded,
2. there exists a saddle-point  $(p', \xi'_n)$  such that  $p^{(k)} \rightarrow p'$  and  $\xi_n^{(k)} \rightarrow \xi'_n$ .

# Implementing a Riemannian manifold



`ManifoldsBase.jl` introduces a manifold type with its field  $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}$  as parameter to provide an interface for implementing functions like

- ▶ `inner(M, p, X, Y)` for the Riemannian metric  $(X, Y)_p$
- ▶ `exp(M, p, X)` and `log(M, p, q)`,
- ▶ more general: `retract(M, p, X, m)`, where `m` is a retraction method
- ▶ similarly: `parallel_transport(M, p, X, q)` and `vector_transport_to(M, p, X, q, m)`

for your manifold, which is a `subtype` of `Manifold`.

😊 mutating version `exp!(M, q, p, X)` works in place in `q`

⊕ basis for generic algorithms working on `any` `Manifold` and generic functions like `norm(M,p,X)`, `geodesic(M, p, X)` and `shortest_geodesic(M, p, q)`

 [juliamanifolds.github.io/ManifoldsBase.jl/](https://github.com/juliamanifolds/ManifoldsBase.jl)



# Manifolds.jl: A Library of Manifolds in Julia

Manifolds.jl is based on the `ManifoldsBase.jl` interface. [Axen, Baran, RB, and Rzecki 2021]

## Features.

- ▶ different metrics
- ▶ Lie groups
- ▶ Build manifolds using
  - ▶ Product manifold  $\mathcal{M}_1 \times \mathcal{M}_2$
  - ▶ Power manifold  $\mathcal{M}^{n \times m}$
  - ▶ Tangent bundle
- ▶ perform statistics
- ▶ well-documented, including formulae and references
- ▶ well-tested, >98 % code coverage, unified coding style

## Manifolds. For example

- ▶ (unit) Sphere
- ▶ Circle & Torus
- ▶ Fixed Rank Matrices
- ▶ Stiefel & Grassmann
- ▶ Hyperbolic space
- ▶ Rotations
- ▶ Symmetric positive definite matrices
- ▶ Symplectic & Symplectic Stiefel
- ▶ ...

 [juliamanifolds.github.io/Manifolds.jl/](https://juliamanifolds.github.io/Manifolds.jl/)



# Manopt.jl: Optimisation on Manifolds in Julia

**Goal.** Provide optimisation algorithms on [Riemannian manifolds](#), i. e. based on [ManifoldsBase.jl](#) such that it can be used with all manifolds from [Manifolds.jl](#).

## Features.

- ▶ generic algorithm framework:  
With `Problem P` and `Options O`
  - ▶ `initialize_solver!(P,O)`
  - ▶ `step_solver!(P, O, i)`: *i*th step
- ➡ run algorithm: call `solve(P,O)`
- ▶ generic debug and recording
- ▶ step sizes and stopping criteria.

## Manopt Family.



[manoptjl.org](https://manoptjl.org)

[RB 2022]



[manopt.org](https://manopt.org) [Boumal, Mishra, Absil, and Sepulchre 2014]



[pymanopt.org](https://pymanopt.org) [Townsend, Koep, and Weichwald 2016]

## Algorithms.

- ▶ Gradient Descent  
CG, Stochastic, Momentum, ...
- ▶ Quasi-Newton  
BFGS, DFP, Broyden, SR1, ...
- ▶ Nelder-Mead, Particle Swarm
- ▶ Subgradient Method
- ▶ Trust Regions
- ▶ Chambolle-Pock
- ▶ Douglas-Rachford
- ▶ Cyclic Proximal Point

# The $\ell^2$ -TV Model

[Rudin, Osher, and Fatemi 1992; Lellmann, Strekalovskiy, Koetter, and Cremers 2013; Weinmann, Demaret, and Storath 2014]

For a manifold-valued image  $f \in \mathcal{M}$ ,  $\mathcal{M} = \mathcal{N}^{d_1, d_2}$ , we compute

$$\arg \min_{p \in \mathcal{M}} \frac{1}{\alpha} F(p) + G(\Lambda(p)), \quad \alpha > 0,$$


with

- ▶ data term  $F(p) = \frac{1}{2} d_{\mathcal{M}}^2(p, f)$
- ▶ “forward differences”  $\Lambda: \mathcal{M} \rightarrow (T\mathcal{M})^{d_1-1, d_2-1, 2}$ ,


$$p \mapsto \Lambda(p) = \left( (\log_{p_i} p_{i+e_1}, \log_{p_i} p_{i+e_2}) \right)_{i \in \{1, \dots, d_1-1\} \times \{1, \dots, d_2-1\}}$$

- ▶ prior  $G(X) = \|X\|_{g,q,1}$  similar to a collaborative TV [Duran, Moeller, Sbert, and Cremers 2016]
- ⇒  $\text{prox}_{\lambda G_n^*}$  given in closed form for  $q = 1$  (anisotropic TV) and  $q = 2$  (isotropic TV).

## Numerical Example for a $\mathcal{P}(3)$ -valued Image




$\mathcal{P}(3)$ -valued data.




anisotropic TV,  $\alpha = 6$ .

- ▶ in each **pixel** we have a symmetric positive definite matrix
- ▶ Applications: denoising/inpainting e.g. of DT-MRI data

# Numerical Example for a $\mathcal{P}(3)$ -valued Image



$\mathcal{P}(3)$ -valued data.




anisotropic TV,  $\alpha = 6$ .

**Approach.** CPPA as benchmark [Bačák 2014; RB, Herzog, Silva Louzeiro, Tenbrinck, and Vidal-Núñez 2021]

	CPPA	PDRA	IRCPA
<b>parameters</b>	$\lambda_k = \frac{4}{k}$	$\lambda = 0.58$ $\beta = 0.93$	$\sigma = \tau = 0.4$ $\gamma = 0.2, m = l$
<b>iterations</b>	4000	122	<b>113</b>
<b>runtime</b>	1235 s.	380 s.	<b>96.1 s.</b>

# Numerical Example for a $\mathcal{P}(3)$ -valued Image




**Approach.** CPPA as benchmark [Bačák 2014; RB, Herzog, Silva Louzeiro, Tenbrinck, and Vidal-Núñez 2021]

	CPPA	PDRA	IRCPA
<b>parameters</b>	$\lambda_k = \frac{4}{k}$	$\lambda = 0.58$ $\beta = 0.93$	$\sigma = \tau = 0.4$ $\gamma = 0.2, m = l$
<b>iterations</b>	4000	122	<b>113</b>
<b>runtime</b>	1235 s.	380 s.	<b>96.1 s.</b>

## Basepoint Effect on $S^2$ -valued Data




Original data



Result,  $m$  west (per px.)

- ▶ piecewise constant results for both
- ! different linearizations lead to different models


# Basepoint Effect on $S^2$ -valued Data



Result,  $m$  mean (per px.)



Result,  $m$  west (per px.)



# Summary

## Summary.

- ▶ We introduced a duality framework on manifolds
- ▶ we introduced a Riemannian Chambolle–Pock algorithm
- ▶ We saw a Software framework for Optimisation algorithms on manifolds
- ▶ Numerical examples illustrates its performance

## Outlook.

- ▶ Strategies for choosing base points, investigate  $C(k)$
- ▶ Investigate constraint optimisation on Manifolds
- ▶ look into further applications

## Selected References

- 
 Axen, S. D., M. Baran, RB, and K. Rzecki (2021). *Manifolds.jl: An Extensible Julia Framework for Data Analysis on Manifolds*. arXiv: 2106.08777.
- 
 Bačák, M. (2014). “Computing medians and means in Hadamard spaces”. In: *SIAM Journal on Optimization* 24.3, pp. 1542–1566. DOI: 10.1137/140953393.
- 
 RB (2022). “Manopt.jl: Optimization on Manifolds in Julia”. In: *Journal of Open Source Software* 7.70, p. 3866. DOI: 10.21105/joss.03866.
- 
 RB, R. Herzog, and M. Silva Louzeiro (2021). *Fenchel duality and a separation theorem on Hadamard manifolds*. arXiv: 2102.11155.
- 
 RB, R. Herzog, M. Silva Louzeiro, D. Tenbrinck, and J. Vidal-Núñez (Jan. 2021). “Fenchel duality theory and a primal-dual algorithm on Riemannian manifolds”. In: *Foundations of Computational Mathematics*. DOI: 10.1007/s10208-020-09486-5. arXiv: 1908.02022.
- 
 RB, J. Persch, and G. Steidl (2016). “A parallel Douglas Rachford algorithm for minimizing ROF-like functionals on images with values in symmetric Hadamard manifolds”. In: *SIAM Journal on Imaging Sciences* 9.4, pp. 901–937. DOI: 10.1137/15M1052858.
- 
 Chambolle, A. and T. Pock (2011). “A first-order primal-dual algorithm for convex problems with applications to imaging”. In: *Journal of Mathematical Imaging and Vision* 40.1, pp. 120–145. DOI: 10.1007/s10851-010-0251-1.